

Corrections for second printing of NUMERICAL METHODS FOR LEAST SQUARES PROBLEMS

Åke Björck

September 10, 2010

Chapter 1

p. 3, l.-16: change “and $\mathcal{V}(z) = \mu \|b\|_2$ ” to “Then if $V = \sigma^2 I$ we have $\mathcal{V}(z) = \sigma^2 f^T f = \sigma^2 \|f\|_2^2$ ”

Chapter 2

- p. 49, l.21: \sqrt{u} should be: $\sqrt{1/u}$
p. 75, l.11,12: change p to r (4 ×)
p. 77, l. 2: should be: $T = L_{21}L_{11}^{-1}, \dots$
p. 87, l. 12: should be: ”is to take ...”
p. 87, l.-12: should be: ”in $B^T B - \tau I$ ”
p. 113, l.-10: should be: $= (V_{22}^T, V_{12}^T).$ ”
p. 113, l.-9: should be: $\Pi = \Pi_I \tilde{\Pi} \Pi_I^T$ ”

Chapter 4

- p. 168, l. -3: first component in b should be: 3
p. 184, l. 18: should be: $|a_i^T x - b_i|^2$.
p. 186, l. 6: should be: $\sigma_1 \geq \sigma_2 \geq 0$

Chapter 5

- p. 188, l.-6: should be: $Ax - b = \hat{A}_2 \bar{x}_2 - \hat{b}$.
p. 189, l. 5: should be: $\hat{A}_2 v = \bar{A}_2 v - \bar{A}_1 R_{11}^{-1} R_{12} v = 0$.
p. 203, l.17: change E_B and $E_{\mathcal{F}}$ to E_B^T and $E_{\mathcal{F}}^T$.

- p. 203, l.20: should be: $\lambda = -U^T(d - UE_{\mathcal{B}}^T x)$.
- p. 205, l.-3. should be: $\psi(x, \lambda) =$
- p. 208, l.15. should be: with $C = I_n$ and
- p. 208, l.-5. should be: is upper bidiagonal
- p. 212, l.-3. should be: $P_\mu = A(A^T A + \mu^2 I_n)^{-1} A^T = \dots$
- p. 213, l. 2 should be: ...
- $$\sum_{i=1}^n \left(\frac{\mu^2 c_i}{\sigma_i^2 + \mu^2} \right)^2 + \dots$$
- p. 213, l. 4 should be: ...
- $$= m - n + \sum_{i=1}^n \frac{\mu^2}{\sigma_i^2 + \mu^2}$$
- p. 213, l. 7 should be: Eldén [272, 1984]

Chapter 6

- p. 217, l.-15: should be: $p = \max_{i \leq i \leq n} (i - f_i(C))$.
- p. 218, l. 16: should be:
 $\text{Env}(C) = \{(i, j) \mid f_i(C) \leq j \leq i\}$.
- p. 225, l. 13–14: $C = \sum_{k=1}^M B_k^T B_k$.

We assume in the following that ...

- p. 225, l. -10– -11: Identifying the blocks in $A^T A = R^T R$, we find that

$$R_i^T R_i = A_i^T A_i, \quad R_i^T S_i = A_i^T B_i, \quad i = 1, \dots, M,$$

$$R_{M+1}^T R_{M+1} = C - \sum_{k=1}^M S_k^T S_k.$$

- p. 246, l.-7: should be: its graph $G(A^T A)$

Chapter 7

- p. 300, l.-10: rhs in (7.5.5) should be: $-C^T(b_2 - Cb_1)$
- p. 309, l. 1: formula should be: $\|r_k\|_2 = \bar{\phi}_{k+1} = \beta_1 s_k s_{k-1} \cdots s_1$.
- p. 311, l. 14: should be: $(\beta_1 e_1, B_k) = P_{k+1} \Omega_k Q_{k+1}$
- p. 311, l. -5: delete the sentence “The SVD of this matrix ...”

Chapter 8

- p. 325, l. 11: should be: matrix $V \in \mathbf{R}^{m \times n}$
- p. 338, l. 1 should be: where $D \in \mathbf{R}^{q \times n}$,
- p. 338, l. 3: should be: $(A \otimes B)d = \text{vec}(BDA^T)$
- p. 338, l. 6: should be $= \text{vec}(B^{-1}DA^{-T})$
- p. 338, l. 9: should be: $= \text{vec}(B^\dagger D(A^\dagger)^T)$

Chapter 9

- p. 343, Example 9.2.1: The point $x^* = 0$ is NOT a local minimizer for $\lambda \geq 1$, but it always is a critical point.
- p. 346, l. -1: change $-$ to $+$ in displayed formula
- p. 346, l.-5: change μ_k to μ_k^2
- p. 347, l. 1, 2 : should read “and for large $\mu_k \dots$ steepest descent direction $-J(x_k)^T r(x_k)$.”
- p. 347, l.-9: should be $\dots - \|r(x_k) + p_k\|_2^2 \dots$
- p. 349, l. 10: should be $(\Sigma_1^2 + V_1^T Q_k V_1)q_1 =$
- p. 349, l.-19: should be maintain p close \dots
- p. 350, l. 17: should be $z_k = J(x_k)^T r(x_k) - J(x_{k-1})^T r(x_k)$
- p. 350, l. 21: change y_k to z_k in equation (9.3.6)
- p. 351, l. -8: should be: based on (9.4.3) \dots
- p. 352, l. -13: should be $B(z)y = \dots$
- p. 356, l. -8: change Q_2 to Q_1 in second formula.
- p. 356, l. -1: change s_2 to s_1 .